

MULTIPLICATIVE FUNCTIONS COMMUTABLE WITH SUMS OF SQUARES

POO-SUNG PARK

ABSTRACT. Let k be an integer greater than or equal 4. We show that if a multiplicative function f satisfies

$$f(x_1^2 + x_2^2 + \cdots + x_k^2) = f(x_1)^2 + f(x_2)^2 + \cdots + f(x_k)^2$$

for all positive integers x_i 's, then f is the identity function.

1. INTRODUCTION

We call an arithmetic function $f(x) : \mathbb{N} \rightarrow \mathbb{C}$ *multiplicative* if $f(1) = 1$ and $f(mn) = f(m)f(n)$ whenever m and n are relatively prime. Many mathematicians have solved various functional equations about multiplicative functions. Claudia Spiro, in 1992, showed that if a multiplicative function f satisfies $f(p+q) = f(p) + f(q)$ for any primes p and q and $f(p_0) \neq 0$ for some prime p_0 , then f is the identity function [15]. It has influenced lots of mathematicians to develop her problem into diverse variations. The number of primes has been varied [9, 7] and primes has been changed to triangular numbers [5]. Lots of articles of similar themes are found in [2, 3, 4, 5, 6, 11, 13, 14].

Recently, Bašić [1] characterized arithmetic functions satisfying $f(m^2 + n^2) = f(m)^2 + f(n)^2$ for all $m, n \in \mathbb{N}$. The author extended his result to the sum of three squares [12]. In this article, the author characterizes multiplicative functions that preserve sum of fixed number of squares.

2. RESULTS

Theorem 1. *If a multiplicative function $f : \mathbb{N} \rightarrow \mathbb{C}$ satisfies*

$$f(x^2 + y^2) = f(x)^2 + f(y)^2$$

for all positive integers x and y , then $f(n^2) = f(n)^2 = n^2$.

Proof. This is a consequence of Bašić's characterization of arithmetic functions satisfying the abovementioned condition [1]. $\square$

Theorem 2. *If a multiplicative function $f : \mathbb{N} \rightarrow \mathbb{C}$ satisfies*

$$f(x^2 + y^2 + z^2) = f(x)^2 + f(y)^2 + f(z)^2$$

for all positive integers x, y and z , then f is the identity function.

Proof. This was proved in [12]. $\square$

Theorem 3. *Let k be an integer ≥ 4 . If a multiplicative function $f : \mathbb{N} \rightarrow \mathbb{C}$ satisfies*

$$f(x_1^2 + x_2^2 + \cdots + x_k^2) = f(x_1)^2 + f(x_2)^2 + \cdots + f(x_k)^2$$

for all positive integers x_i , then f is the identity function.

Proof. Dubouis [8], [10, Chapter 6] showed that every integer n can be represented as a sum of k nonvanishing squares except

$$n = \begin{cases} 1, 3, 5, 9, 11, 17, 29, 41, 2 \cdot 4^m, 6 \cdot 4^m, 14 \cdot 4^m \ (m \geq 0) & \text{if } k = 4, \\ 33 & \text{if } k = 5, \\ 1, 2, \dots, k-1, k+1, k+2, k+4, k+5, k+7, k+10, k+13 & \text{if } k \geq 5. \end{cases}$$

i) Assume that $k = 4$. Note that $f(1) = 1$ and $f(4) = 4$.

Since

$$\begin{aligned} f(12) &= f(3)f(4) = 4f(3) \\ &= f(1)^2 + f(1)^2 + f(1)^2 + f(3)^2 = 3 + f(3)^2, \end{aligned}$$

we obtain that $f(3) = 1$ or $f(3) = 3$.

Let us consider $f(5)$. Since

$$\begin{aligned} f(20) &= f(4)f(5) = 4f(5) \\ &= f(1)^2 + f(1)^2 + f(3)^2 + f(3)^2 = 2 + 2f(3)^2, \end{aligned}$$

$f(5) = 1$ or $f(5) = 5$ according to the value of $f(3)$.

Similarly, from

$$\begin{aligned} f(28) &= f(4)f(7) = 4f(7) \\ &= f(1)^2 + f(3)^2 + f(3)^2 + f(3)^2 = 1 + 3f(3)^2 \end{aligned}$$

we have that $f(7) = 1$ or $f(7) = 7$ according to the value of $f(3)$.

Note that

$$\begin{aligned} f(35) &= f(5)f(7) \\ &= f(1)^2 + f(3)^2 + f(3)^2 + f(4)^2 = 17 + 2f(3)^2. \end{aligned}$$

Hence we conclude that $f(3) = 3$, $f(5) = 5$, and $f(7) = 7$. Then $f(2) = 2$ also follows from

$$\begin{aligned} f(10) &= f(2)f(5) \\ &= f(1)^2 + f(1)^2 + f(2)^2 + f(2)^2 = 2 + 2f(2)^2 \end{aligned}$$

and

$$f(7) = f(1)^2 + f(1)^2 + f(1)^2 + f(2)^2 = 3 + f(2)^2.$$

If we consider

$$\begin{aligned} f(18) &= f(2)f(9) = 2f(9) \\ &= f(1)^2 + f(2)^2 + f(2)^2 + f(3)^2 = 18, \end{aligned}$$

we have that $f(9) = 9$. The similar manner gives us

$$f(11) = 11, \quad f(17) = 17, \quad f(29) = 29, \quad f(41) = 41.$$

Now suppose that $f(2 \cdot 4^r) = 2 \cdot 4^r$ for $r < m$. Then $f(n) = n$ when $n < 2 \cdot 4^m$. Since $5 \cdot 2 \cdot 4^m$ can be represented as a sum of squares of four positive integers less than $2 \cdot 4^m$, we can conclude that $f(5 \cdot 2 \cdot 4^m) = 5 \cdot 2 \cdot 4^m$. In other hand,

$$f(5 \cdot 2 \cdot 4^m) = f(5)f(2 \cdot 4^m) = 5f(2 \cdot 4^m)$$

and thus $f(2 \cdot 4^m) = 2 \cdot 4^m$.

Now consider $6 \cdot 4^m$. Since $f(6 \cdot 4^m) = f(3)f(2 \cdot 4^m)$, we obtain that $f(6 \cdot 4^m) = 6 \cdot 4^m$. Similarly, $f(14 \cdot 4^m) = f(7)f(2 \cdot 4^m) = 14 \cdot 4^m$.

ii) Assume $k = 5$. Then $f(1) = 1$ and $f(5) = 5$.

From

$$\begin{aligned} f(20) &= f(4)f(5) = 5f(4) \\ &= f(1)^2 + f(1)^2 + f(1)^2 + f(1)^2 + f(4)^2 = 4 + f(4)^2 \\ &= f(2)^2 + f(2)^2 + f(2)^2 + f(2)^2 + f(2)^2 = 5f(2)^2 \end{aligned}$$

we obtain $f(4) = 1$ or $f(4) = 4$. Also, $f(2) = 1$ or $f(2) = \pm 2$ according to the value of $f(4)$.

From

$$\begin{aligned} f(29) &= f(1)^2 + f(1)^2 + f(1)^2 + f(1)^2 + f(5)^2 = 29 \\ &= f(1)^2 + f(1)^2 + f(3)^2 + f(3)^2 + f(3)^2 = 2 + 3f(3)^2 \\ &= f(1)^2 + f(2)^2 + f(2)^2 + f(2)^2 + f(4)^2 = 1 + 3f(2)^2 + f(4)^2 \end{aligned}$$

we obtain that $f(29) = 29$, $f(2) = \pm 2$, and $f(3) = \pm 3$. Also, this fixes $f(4) = 4$.

Note that both of m and $2m$ can be represented as sums of five squares for sufficiently large m coprime to 2. Since $f(2m) = f(2)f(m)$ and both of $f(2m)$ and $f(m)$ are positive, $f(2)$ is also positive. That is, $f(2) = 2$. Similarly, $f(3) = 3$ and thus $f(6) = 6$.

Now, we use induction to show that $f(n) = n$ for $n \geq 7$. Assume that $f(m) = m$ for $m < n$. Then $n(n-1) \geq 42$ and it can be represented as a sum of squares of five positive integers less than n . Thus

$$f(n(n-1)) = n(n-1).$$

Since n and $n-1$ are relatively prime,

$$f(n(n-1)) = f(n)f(n-1) = f(n)(n-1)$$

and thus $f(n) = n$.

iii) $k = 6$

Clearly, $f(1) = 1$ and $f(6) = 6$. From the equalities

$$\begin{aligned} f(30) &= f(5)f(6) = 6f(5) \\ &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2) = 5 + f(5)^2 \\ &= f(1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2) = 3 + 3f(3)^2 \\ &= f(1^2 + 1^2 + 2^2 + 2^2 + 2^2 + 4^2) = 2 + 3f(2)^2 + f(4)^2 \\ f(41) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 6^2) = 41 \\ &= f(1^2 + 1^2 + 1^2 + 2^2 + 3^2 + 5^2) = 3 + f(2)^2 + f(3)^2 + f(5)^2 \\ f(21) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 4^2) = 5 + f(4)^2 \\ &= f(1^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2) = 1 + 5f(2)^2 \end{aligned}$$

we deduce that

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(4) = \pm 4, \quad f(5) = 5.$$

Considering $2m$, $3m$, and $4m$ for sufficiently large m , we obtain that $f(n) = n$ for $n \leq 6$. Now we can show $f(n) = n$ by the way similar to the previous case.

iv) $k = 7$

We have that $f(1) = 1$ and $f(7) = 7$. Note that

$$\begin{aligned}
f(31) &= f(1^2 + 1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2) = 4 + 3f(3)^2 \\
&= f(1^2 + 1^2 + 1^2 + 2^2 + 2^2 + 2^2 + 4^2) = 3 + 3f(2)^2 + f(4)^2 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2) = 6 + f(5)^2 \\
f(42) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 2^2 3^2) = 6 + f(2)^2 f(3)^2 \\
&= f(1^2 + 2^2 + 2^2 + 2^2 + 2^2 + 3^2 + 4^2) = 1 + 4f(2)^2 + f(3)^2 + f(4)^2 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 2^2 + 3^2 + 5^2) = 4 + f(2)^2 + f(3)^2 + f(5)^2 \\
f(55) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 7^2) = 55 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2 + 5^2) = 5 + 2f(5)^2.
\end{aligned}$$

Thus, we can find that

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(4) = \pm 4, \quad f(5) = \pm 5.$$

We can conclude that $f(n) = n$ by the similar way.

v) $k \geq 8$

Since $f(1) = 1$, $f(k) = k$, and

$$f(k(k-1)) = f(k)f(k-1) = \underbrace{f(1)^2 + \cdots + f(1)^2}_{k-1 \text{ times}} + f(k-1)^2,$$

the value of $f(k-1)$ is either 1 or $k-1$.

Since

$$\begin{aligned}
40 &= 1^2 + 1^2 + 1^2 + 1^2 + 6^2 \\
&= 2^2 + 3^2 + 3^2 + 3^2 + 3^2,
\end{aligned}$$

we obtain that

$$(k-5) \times f(1)^2 + 4 \times f(1)^2 + (f(2)f(3))^2 = (k-5) \times f(1)^2 + f(2)^2 + 4f(3)^2$$

or

$$4 + (f(2)f(3))^2 = f(2)^2 + 4f(3)^2.$$

Also, from

$$\begin{aligned}
32 &= 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2 \\
&= 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2,
\end{aligned}$$

we obtain that $5 + 3f(3)^2 = 8f(2)^2$.

Solving two equations, we can find solutions

$$f(2) = \pm 1, f(3) = \pm 1 \quad \text{or} \quad f(2) = \pm 2, f(3) = \pm 3.$$

Now, let $3a + 8b = 2k - 1$ with nonnegative integers a and b . This is possible unless $2k - 1 = 1, 2, 4, 5, 7, 10, 13$. That is, if $2k - 1 > 13$, then it is represented in such a way. The number 13 is called the *Frobenius number* of 3 and 8.

Note that

$$(k-1) + (2k-1) + (k-1)^2 = k^2 + k - 1$$

and thus

$$\begin{aligned}
 f(k^2 + k - 1) &= f(k)^2 + (k - 1)f(1)^2 = k^2 + k - 1 \\
 &= f(\underbrace{1^2 + \cdots + 1^2}_{k-a-b-1 \text{ times}} + \underbrace{2^2 + \cdots + 2^2}_a + \underbrace{3^2 + \cdots + 3^2}_b + (k - 1)^2) \\
 &= (k - a - b - 1) + af(2)^2 + bf(3)^2 + f(k - 1)^2.
 \end{aligned}$$

Therefore

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(k - 1) = k - 1.$$

Since

$$\begin{aligned}
 4^2 + 2^2 + 2^2 + 2^2 &= 1^2 + 3^2 + 3^2 + 3^2 \\
 5^2 + 1^2 + 1^2 &= 3^2 + 3^2 + 3^2 \\
 6 &= 2 \times 3 \\
 7^2 + 1^2 &= 5^2 + 5^2 \\
 8^2 + 1^2 &= 4^2 + 7^2 \\
 9^2 + 2^2 &= 6^2 + 7^2 \\
 10 &= 2 \times 5,
 \end{aligned}$$

we obtain that $f(n) = \pm n$ for $n \leq 10$.

Note that

$$(2\ell + 1)^2 + (\ell - 2)^2 = (2\ell - 1)^2 + (\ell + 2)^2 \text{ with } \ell \geq 5$$

and

$$(2\ell)^2 + (\ell - 5)^2 = (2\ell - 4)^2 + (\ell + 3)^2 \text{ with } \ell \geq 6.$$

Hence, by induction, $f(n) = \pm n$ for all n .

It is similar to the previous cases to show that $f(n)$ is positive. Therefore, $f(n) = n$.

□

REFERENCES

- [1] Bašić, B.; Characterization of arithmetic functions that preserve the sum-of-squares operation. *Acta Math. Sin.*, **30** (2014), 689–695.
- [2] Chen, K.-K.; Chen, Y.-G.; On $f(p) + f(q) = f(p + q)$ for all odd primes p and q . *Publ. Math. Debrecen*, **76** (2010), 425–430.
- [3] Chen, Y.-G.; Fang, J.-H.; Yuan, P.; Zheng, Y.; On multiplicative functions with $f(p + q + n_0) = f(p) + f(q) + f(n_0)$. *J. Number Theory*, **165** (2016), 270–289.
- [4] Chung, P. V.; Multiplicative functions satisfying the equation $f(m^2 + n^2) = f(m^2) + f(n^2)$. *Math. Slovaca*, **46** (1996), 165–171.
- [5] Chung, P. V.; Phong, B. M.; Additive uniqueness sets for multiplicative functions. *Publ. Math. Debrecen*, **55** (1999), 237–243.
- [6] De Koninck, J.-M.; Kátai, I.; Phong, B. M.; A new characteristic of the identity function. *J. Number Theory*, **63** (1997), 325–338.
- [7] Dubickas, A.; Šarka, P.; On multiplicative functions which are additive on sums of primes. *Aequat. Math.*, **86** (2013), 81–89.
- [8] Dubouis, E.; Solution of a problem of J. Tannery, *L'Intermédiaire des Math.*, **18** (1911), 55–56.
- [9] Fang, J.-H.; A characterization of the identity function with equation $f(p + q + r) = f(p) + f(q) + f(r)$. *Combinatorica*, **31** (2011), 697–701.
- [10] Grosswald, E.; Representations of Integers as Sums of Squares. Springer, 2011.

- [11] Indlekofer, K.-H.; Phong, B. M.; Additive uniqueness sets for multiplicative functions. *Ann. Univ. Sci. Budapest. Sect. Comput.*, **26** (2006), 65–77.
- [12] Park, P.-S.; Multiplicative functions preserving the sum-of-three-squares operation. submitted.
- [13] Phong, B. M.; On sets characterizing the identity function. *Ann. Univ. Sci. Budapest. Sect. Comput.*, **24** (2004), 295–306.
- [14] Phong, B. M.; A characterization of the identity function with an equation of Hosszú type. *Publ. Math. Debrecen*, **69** (2006), 219–226.
- [15] Spiro, C. A.; Additive uniqueness sets for arithmetic functions. *J. Number Theory*, 42 (1992), 232–246.